Complete Summary

TITLE

Eye care: percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy AND the presence or absence of macular edema during one or more office visits within 12 months.

SOURCE(S)

American Academy of Ophthalmology, Physician Consortium for Performance Improvement®, National Committee for Quality Assurance. Eye care physician performance measurement set. Chicago (IL): American Medical Association, National Committee for Quality Assurance; 2007 Oct. 13 p. [8 references]

Measure Domain

PRIMARY MEASURE DOMAIN

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

SECONDARY MEASURE DOMAIN

Does not apply to this measure

Brief Abstract

DESCRIPTION

This measure is used to assess the percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy AND the presence or absence of macular edema during one or more office visits within 12 months.

RATIONALE

Several level 1 randomized control trial (RCT) studies demonstrate the ability of timely treatment to reduce the rate and severity of vision loss from diabetes (Diabetic Retinopathy Study-DRS, Early Treatment Diabetic Retinopathy Study-

ETDRS). Necessary examination prerequisites to applying the study results are that the presence and severity of both peripheral diabetic retinopathy and macular edema be accurately documented. In the RAND chronic disease quality project, while administrative data indicated that roughly half of the patients had an eye exam in the recommended time period, chart review data indicated that only 19% had documented evidence of a dilated examination. Thus, ensuring timely treatment that could prevent 95% of the blindness due to diabetes requires the performance and documentation of key examination parameters. The documented level of severity of retinopathy and the documented presence or absence of macular edema assists with the on-going plan of care for the patient with diabetic retinopathy.*

*The following clinical recommendation statement is quoted <u>verbatim</u> from the referenced clinical guidelines and represents the evidence base for the measure:

Since treatment is effective in reducing the risk of visual loss, detailed examination is indicated to assess for the following features that often lead to visual impairment: presence of macular edema, optic nerve neovascularization and/or neovascularization elsewhere, signs of severe nonproliferative diabetic retinopathy (NPDR) and vitreous or preretinal hemorrhage. (American Academy of Ophthalmology [AAO])

PRIMARY CLINICAL COMPONENT

Diabetic retinopathy; dilated macular or fundus exam; documentation of level of severity of retinopathy; documentation of presence or absence of macular edema

DENOMINATOR DESCRIPTION

All patients aged 18 years and older with a diagnosis of diabetic retinopathy (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary)

NUMERATOR DESCRIPTION

Patients who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy AND the presence or absence of macular edema during one or more office visits within 12 months (see the related "Numerator Inclusions/Exclusions" field in the Complete Summary)

Evidence Supporting the Measure

EVIDENCE SUPPORTING THE CRITERION OF QUALITY

- A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence
- One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

NATIONAL GUIDELINE CLEARINGHOUSE LINK

Diabetic retinopathy.

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Use of this measure to improve performance

EVIDENCE SUPPORTING NEED FOR THE MEASURE

McGlynn EA, Asch SM, Adams J, Keesey J, Hicks J, DeCristofaro A, Kerr EA. The quality of health care delivered to adults in the United States. N Engl J Med2003 Jun 26;348(26):2635-45. PubMed

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Internal quality improvement National reporting

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care Physician Group Practices/Clinics

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Age greater than or equal to 18 years

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Unspecified

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Unspecified

UTILIZATION

Unspecified

COSTS

Unspecified

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

All patients aged 18 years and older with a diagnosis of diabetic retinopathy

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients aged 18 years and older with a diagnosis of diabetic retinopathy

Exclusions

- Documentation of medical reason(s) for not performing a dilated macular or fundus examination
- Documentation of patient reason(s) for not receiving a dilated macular or fundus examination
- Documentation of system reason(s) for not receiving a dilated macular or fundus examination

Note: The system reason exclusion may be used if a physician is asked to report on this measure but is not the physician providing the primary management for diabetic retinopathy.

RELATIONSHIP OF DENOMINATOR TO NUMERATOR

All cases in the denominator are equally eligible to appear in the numerator

DENOMINATOR (INDEX) EVENT

Clinical Condition Encounter

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Patients who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy AND the presence or absence of macular edema during one or more office visits within 12 months

Note: Medical record must include:

• Documentation of the level of severity of retinopathy (e.g., background diabetic retinopathy, proliferative diabetic retinopathy, nonproliferative diabetic retinopathy)

AND

Documentation of whether macular edema was present or absent.

Exclusions

None

MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

NUMERATOR TIME WINDOW

Encounter or point in time

DATA SOURCE

Administrative data Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Measure #7 diabetic retinopathy: documentation of presence or absence of macular edema and level of severity of retinopathy.

MEASURE COLLECTION

The Physician Consortium for Performance Improvement® Measurement Sets

MEASURE SET NAME

Eye Care Physician Performance Measurement Set

SUBMITTER

American Medical Association on behalf of the American Academy of Ophthalmology, the National Committee for Quality Assurance, and the Physician Consortium for Performance Improvement®

DEVELOPER

American Academy of Ophthalmology National Committee for Quality Assurance Physician Consortium for Performance Improvement®

FUNDING SOURCE(S)

Unspecified

COMPOSITION OF THE GROUP THAT DEVELOPED THE MEASURE

Paul P. Lee, MD, JD (*Co-Chair*); Jinnet B. Fowles, PhD (*Co-Chair*); Richard L. Abbott, MD; Lloyd P. Aiello, MD, PhD; Priscilla P. Arnold, MD; Richard Hellman, MD, FACP, FACE; Leon W. Herndon, MD; Kenneth J. Hoffer, MD; Jeffrey S. Karlik, MD; Mathew W. MacCumber, MD, PhD; Mildred M.G. Olivier, MD; James L. Rosenzweig, MD, FACE; Sam J.W. Romeo, MD, MBA; John T. Thompson, MD

Andrea Gelzer, MD, MS, FACP, Health Plan Representative (HealthNet)

Rebecca A. Kresowik, Consortium Consultant; Timothy F. Kresowik, MD, Consortium Consultant

Flora Lum, MD, American Academy of Ophthalmology

Heidi Bossley, MSN, MBA, American Medical Association; Karen Kmetik, PhD, American Medical Association

FINANCIAL DISCLOSURES/OTHER POTENTIAL CONFLICTS OF INTEREST

Conflicts, if any, are disclosed in accordance with the Physician Consortium for Performance Improvement® conflict of interest policy.

ENDORSER

National Quality Forum

INCLUDED IN

Ambulatory Care Quality Alliance Physician Quality Reporting Initiative

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2006 Oct

REVISION DATE

2007 Oct

MEASURE STATUS

This is the current release of the measure.

SOURCE(S)

American Academy of Ophthalmology, Physician Consortium for Performance Improvement®, National Committee for Quality Assurance. Eye care physician performance measurement set. Chicago (IL): American Medical Association, National Committee for Quality Assurance; 2007 Oct. 13 p. [8 references]

MEASURE AVAILABILITY

The individual measure, "Measure #7 Diabetic Retinopathy: Documentation of Presence or Absence of Macular Edema and Level of Severity of Retinopathy," is published in the "Eye Care Physician Performance Measurement Set." This document and technical specifications are available in Portable Document Format (PDF) from the American Medical Association (AMA)-convened Physician Consortium for Performance Improvement® Web site: www.physicianconsortium.org.

For further information, please contact AMA staff by e-mail at cgi@ama-assn.org.

NQMC STATUS

This NQMC summary was completed by ECRI Institute on February 11, 2008. The information was verified by the measure developer on April 14, 2008.

COPYRIGHT STATEMENT

© Measures including specifications © 2007 American Medical Association and National Committee for Quality Assurance.

CPT® Copyright 2006 American Medical Association

Disclaimer

NQMC DISCLAIMER

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at

http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

© 2008 National Quality Measures Clearinghouse

Date Modified: 11/3/2008

